

# task\_1.2.5\_with\_calc

## Student Group

First Name	Surname	Matrikel Nr.

## Table of Contents

Task 1.2.5 Forces on Charges (exam task, ca 8 % of a 60 minute exam, WS2020) ..... 2

**Task 1.2.5 Forces on Charges (exam task, ca 8 % of a 60 minute exam, WS2020)**



Given is an arrangement of electric charges located in a vacuum (see picture on the right).

The charges have the following values:

$Q_1 = 7 \mu\text{C}$  (point charge)

$Q_2 = 5 \mu\text{C}$  (point charge)

$Q_3 = 0 \text{ C}$  (infinitely extended surface charge)

$\epsilon_0 = 8.854 \cdot 10^{-12} \text{ F/m}$ ,  $\epsilon_r = 1$

1. calculate the magnitude of the force of  $Q_2$  on  $Q_1$ , without the force effect of  $Q_3$ .

Tips for the solution

- Which equation is to be used for the force effect of charges?
- How can the distance between the two charges be determined?

Solution

$$\begin{aligned} F_C &= \frac{1}{4\pi \cdot \epsilon_0} \cdot \frac{Q_1 \cdot Q_2}{r^2} \quad \& \quad | \quad \text{with } r = \sqrt{\Delta x^2 + \Delta y^2} \quad \& \quad F_C = \frac{1}{4\pi \cdot \epsilon_0} \cdot \frac{Q_1 \cdot Q_2}{\Delta x^2 + \Delta y^2} \quad \& \quad | \quad \text{Insert numerical values, read off distances: } \Delta x = 5 \text{ dm}, \Delta y = 3 \text{ dm} \\ F_C &= \frac{1}{4\pi \cdot 8.854 \cdot 10^{-12} \text{ F/m}} \cdot \frac{7 \cdot 10^{-6} \text{ C} \cdot 5 \cdot 10^{-6} \text{ C}}{(0.5 \text{ m})^2 + (0.2 \text{ m})^2} \end{aligned}$$

## Result

$$\begin{align*} |\vec{F}_C| = 1.084 \text{ ~}\mathrm{N} \end{align*} \rightarrow 1.1 \text{ ~}\mathrm{N}$$

2. is this force attractive or repulsive?

## Tips for the solution

- What force effect do equally or oppositely charged bodies exhibit on each other?

## Solution

The force is repulsive because both charges have the same sign.

Now let  $Q_2=0$  and the surface charge  $Q_3$  be designed in such a way that a homogeneous electric field with  $E_3=100 \text{ ~}\mathrm{kV/m}$  results. What force (magnitude) now results on  $Q_1$ ?

## Tips for the solution

- Which equation is to be applied for the force action in the homogeneous field?

## Solution

$$\begin{align*} F_C &= E \cdot Q_1 \quad \& \quad | \text{\textit{Insert numerical values}} \\ F_C &= 100 \text{ ~}\mathrm{V/m} \cdot 7 \cdot 10^{-6} \text{ ~}\mathrm{C} \end{align*}$$

## Result

$$\begin{align*} |\vec{F}_C| = 0.7 \text{ ~}\mathrm{N} \end{align*}$$

From:

<https://mexle.te.hs-heilbronn.de/> - MEXLE Wiki

Permanent link:

[https://mexle.te.hs-heilbronn.de/electrical\\_engineering\\_and\\_electronics\\_1/task\\_1.2.5\\_with\\_calc](https://mexle.te.hs-heilbronn.de/electrical_engineering_and_electronics_1/task_1.2.5_with_calc)

Last update: **2025/10/20 02:45**

