

task_I9hubowt6x00b2h5_with_calculation

Student Group

First Name	Surname	Matrikel Nr.

Table of Contents

Exercise E8 Determining the Current from Charge per Time	2
--	---

current, charge, chapter1 4

Exercise E8 Determining the Current from Charge per Time

Two objects have different charge increases per time duration, figure 1 shows a non-linear increase in the charge per time.

Result



Start drawing by clicking here.



Start drawing by clicking here.

A non-linear charge increase leads to a non-constant current.

For a non-constant current, one has to use the time derivative of the charge Q to get the current I .

So, the formula $I = \frac{dQ}{dt}$ has to be used instead of $I =$

.. Determine the currents I_1 and I_2 for the two objects from the Q - t -diagram

figure 1 and plot the currents into a new diagram.

Solution

- Have a look how much increase ΔQ per time duration Δt is there for each object.
- For this choose a distinct time period, e.g. between 0 s and 20 s .
- The current is then given as the change in charge per time: $I = \frac{\Delta Q}{\Delta t}$

From:

<https://mexle.te.hs-heilbronn.de/> - MEXLE Wiki

Permanent link:

https://mexle.te.hs-heilbronn.de/electrical_engineering_and_electronics/task_I9hubowt6x00b2h5_with_calculation

Last update: 2023/10/03 19:28

